

# FACT SHEET



## COMBINATION POUR-ON FOR CATTLE

**Active Constituents:** ABAMECTIN 10 mg/mL

LEVAMISOLE 200 mg/mL

**Solvents:** 250 mg/mL Diethylene glycol monobutyl ether  
540 mg/mL N-methyl-2-pyrrolidone



## ECLIPSE – Twice the Pour-On.

### PRODUCT DESCRIPTION\*

For the treatment and control of roundworms, including macrocyclic lactone or levamisole resistant strains, and external parasites in cattle.

### PRODUCT BENEFITS

- ✓ Combined power of 2 actives together
- ✓ Protects herds against resistance<sup>^</sup>
- ✓ The only pour-on with a registered claim to control resistant worms
- ✓ Unsurpassed control over *Cooperia* and *Ostertagia* species
- ✓ Convenient low 1 mL per 20 kg dose rate
- ✓ The power of a combination in a pour-on

### BROAD-SPECTRUM ACTIVITY AGAINST RESISTANCE

ECLIPSE is Australia's first dual action broad-spectrum pour-on for cattle.

ECLIPSE contains abamectin, a member of the macrocyclic lactones family of chemicals, which is effective against sensitive strains of internal and external parasites. It also contains levamisole, a member of the imidazothiazoles family of chemicals, which is effective against roundworms, including lungworm.

Being a dual action pour-on, the abamectin component works to kill a broad spectrum of internal and external parasites, while the levamisole component works to kill ML-resistant strains of *Cooperia* and *Ostertagia*. This results in superior, longer-term and sustainable control.

By combining two powerful actives in one single convenient dose, you now have the power of a combination in a pour-on.

### INDICATIONS\*

For the treatment and control of adult and immature stages of:

**Gastrointestinal roundworms, including strains resistant to macrocyclic lactones ("mectin drenches") or to imidazothiazoles ("clear drenches" such as levamisole)**

Barbers Pole Worm (*Haemonchus placei*)  
Small Brown Stomach Worm (*Ostertagia ostertagi*, *O. lyrata*, including inhibited stages)  
Stomach Hair Worm (*Trichostrongylus axei*)  
Small Intestinal Worm (*Cooperia oncophora*, *C. punctata*, *C. pectinata*, *C. mcmasteri*)  
Hookworm (*Bunostomum phlebotomum*)  
Thin-necked Intestinal Worm (*Nematodirus spp.*)  
Intestinal Threadworm (*Strongyloides papilliferus*)  
Black Scour Worm (*Trichostrongylus spp.*)  
Large Bowel Worm (*Oesophagostomum radiatum*)

Large Mouth Bowel Worm (*Chabertia ovina*)  
Whipworm (*Trichuris spp.*)

**Lungworms** (*Dictyocaulus viviparus*)

**Lice**  
Long-nosed Sucking Lice (*Linognathus vituli*)  
Biting Lice (*Bovicola (Damalinia) bovis*)

**Buffalo Fly** (*Haematobia irritans exigua*)

ECLIPSE Pour-On controls infection with Buffalo Fly for up to 10 days after treatment.

**Cattle Tick** (*Boophilus microplus*)

Maximum tick efficacy is achieved 4-5 days after treatment. Engorged female ticks that drop from cattle in the initial days after treatment may lay viable eggs. This should be taken into account when planning a strategic tick control program.



## DIRECTIONS FOR USE

### Dosage and Administration

ECLIPSE Pour-On dose rate is 1 mL / 20 kg body weight (0.5 mg/kg abamectin and 10 mg/kg levamisole). Apply along the backline of the animal, using the ECLIPSE Pour-On applicator, in a strip starting between the shoulder blades.

Only use the pour-on applicator supplied for use with ECLIPSE Pour-On. ECLIPSE is not compatible with other guns. After use, rinse gun with warm soapy water.

**Re-treatment interval:** DO NOT re-treat animals for 35 days after last treatment

### WITHHOLDING PERIODS

**MEAT:** DO NOT USE less than 56 days before slaughter for human consumption.

Calves MUST NOT be slaughtered for human consumption within 60 days of treatment of dams.

**MILK:** DO NOT USE in lactating cows or within 60 days of calving where milk may be used or processed for human consumption. If cows calve earlier than 60 days after treatment, milk may contain residues. This milk MUST NOT be used for human consumption, supplied for processing or fed to bobby calves for at least 60 days following treatment.

**EXPORT SLAUGHTER INTERVAL (ESI)†:** DO NOT USE less than 70 days before slaughter for export.

### RAINFALL

ECLIPSE Pour-On is rainfast. Rain occurring as early as 2 hours after treatment is unlikely to affect the efficacy of the product. However rainfall on cattle less than 2 hours after treatment may result in reduced efficacy. ECLIPSE Pour-On should not be applied to wet animals.

### STORAGE

Store below 25°C (Air Conditioning). Protect from light.

### PACK SIZE

2.5 L back pack & 5 L back pack

Live weight	Dose Volume mL	2.5 Litre pack treats	5 Litre pack treats
51-100kg	0.5 mL / 10 kg	at least 500	at least 1000
101-150kg	7.5	333	666
151-200kg	10	250	500
201-250kg	12.5	200	400
251-300kg	15	166	333
301-350kg	17.5	142	285
351-400kg	20	125	250
401-450kg	22.5	111	222
451-500kg	25	100	200
501-550kg	27.5	90	181
551-600kg	30	83	166
601-650kg	32.5	76	153

DO NOT USE in calves under 50 kg bodyweight.

Over 650 kg, dose at rate of 1 mL/20 kg bodyweight.

Contact: your local Merial Territory Manager, your local rural store, call Merial Customer Solutions on 1800 808 691 or visit [www.merial.com.au](http://www.merial.com.au)

\*See product label for full details.

†The usage of effective multiple active drenches to slow the development of parasite resistance is well supported by peer-review publications and technical opinion.

‡The ESI was correct at the time of publication. Before using this product confirm the current ESI from Merial (1800 808 691) or the APVMA website ([www.apvma.gov.au/residues](http://www.apvma.gov.au/residues)).

Merial Australia Pty Ltd, Building D, 12-24 Talavera Road, Macquarie Park, NSW 2113  
(ABN 53 071 187 285). \*ECLIPSE is a registered trademark of Merial Limited.  
©Merial Limited 2014. All rights reserved. ECLP.14.08.0121



**eclipse®**  
COMBINATION POUR-ON FOR CATTLE

