

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS
FOR ANIMAL TREATMENT ONLY

eclipse®

COMBINATION POUR-ON FOR CATTLE



Active Constituents:
200 mg/mL LEVAMISOLE
10 mg/mL ABAMECTIN

Solvents:
540 mg/mL N-methyl-2-pyrrolidone
250 mg/mL Diethylene glycol monobutyl ether

**FOR THE TREATMENT AND CONTROL OF ROUNDWORMS, INCLUDING MACROCYCLIC LACTONE,
OR LEVAMISOLE RESISTANT STRAINS, AND EXTERNAL PARASITES IN CATTLE**

©ECLIPSE and cattle head logo are registered trademarks of Merial Limited. ©2014 Merial Limited. All rights reserved.

5 Litres



ECLIPSE POUR-ON

ECLIPSE Pour-On contains abamectin, a member of the macrocyclic lactone family of chemicals. It is effective against sensitive strains of internal and external parasites. It also contains levamisole, a member of the imidazothiazole family of chemicals, which is effective against roundworms, including lungworm. Resistance may develop to any chemical.

INDICATIONS

For the treatment and control of adult and immature stages of:

Gastro-intestinal roundworms, including strains resistant to macrocyclic lactones ("mectin drenches") or to imidazothiazoles ("clear drenches" such as levamisole)
Barber's Pole Worm (*Haemonchus placei*), Small Brown Stomach Worm (*Ostertagia ostertagi*, *O. lyrata*, including inhibited stages), Stomach Hair Worm (*Trichostrongylus axei*), Small Intestinal Worm (*Cooperia oncophora*, *C. punctata*, *C. pectinata*, *C. mcmasteri*), Hookworm (*Bunostomum phlebotomum*), Thin-necked Intestinal Worm (*Nematodirus* spp.), Intestinal Threadworm (*Strongyloides papillosus*), Black Scour Worm (*Trichostrongylus* spp.), Large Bowel Worm (*Oesophagostomum radiatum*), Large Mouth Bowel Worm (*Chabertia ovina*), Whipworm (*Trichuris* spp.), **Lungworm** (*Dictyoacaulus viviparus*), **Sucking Lice** (*Linognathus vituli*) **Biting Lice** (*Bovicola* [*Damalinia*] *bovis*), **Buffalo Fly** (*Haematobia irritans exigua*)
ECLIPSE Pour-On controls infection with Buffalo Fly for up to 10 days after treatment.

Cattle Tick (*Boophilus microplus*)
Maximum tick efficacy is achieved 4–5 days after treatment. Engorged female ticks that drop from cattle in the initial days after treatment may lay viable eggs. This should be taken into account when planning a strategic tick control program. ECLIPSE Pour-On should be alternated with other dips or pour-ons when the program requires repeat treatments.

DIRECTIONS FOR USE

Restraints
DO NOT USE in lactating cows or within 60 days of calving where milk may be used or processed for human consumption.
DO NOT USE in calves under 50 kg bodyweight.

Precautions

This product should not be used on animals which are severely stressed or in ill health. Care should be taken when treating pregnant animals to minimise handling stress. Recommended dose rates should not be exceeded. Doses of 3 or more times the recommended dose can cause symptoms of levamisole toxicity, so estimate weights carefully. Dehydrated animals may be more susceptible to toxicity.

Side Effects

Temporary skin reaction or hair loss may occur along the application site, especially if applied during very hot weather. Therefore do not treat cattle shortly prior to showing. These reactions should resolve without treatment. If signs persist or worsen, contact your veterinarian.

Dosage and Administration

ECLIPSE Pour-On dose rate is 1 mL / 20 kg body weight (0.5 mg/kg abamectin and 10 mg/kg levamisole). Apply along the backline of the animal, using the ECLIPSE Pour-On applicator, in a strip starting between the shoulder blades.

This product is not to be given orally or injected. Only use the pour-on applicator supplied for use with ECLIPSE Pour-On. ECLIPSE is not compatible with other guns. After use, rinse gun with warm soapy water.

Re-treatment interval: DO NOT re-treat animals for 35 days after last treatment.

Live weight kg	Dose Volume mL / 10 kg	5 litre pack treats at least 1000
51–100	0.5 mL / 10 kg	at least 1000
101–150	7.5	666
151–200	10	500
201–250	12.5	400
251–300	15	333
301–350	17.5	285
351–400	20	250
401–450	22.5	222
451–500	25	200
501–550	27.5	181
551–600	30	166
601–650	32.5	153

Under 100 kg, dose at rate of 0.5 mL / 10 kg bodyweight, and over 650 kg dose at rate of 1 mL / 20 kg bodyweight. A representative sample of animals should be weighed either with scales or a weighband before treatment. Dose according to the heaviest animal by live weight in the group (cows, bulls, heifers, etc). Do not underdose. Where there is a large variation in size within the group, draft into two or more lines based on bodyweight to avoid excessive overdosing. Check the accuracy of guns regularly.

ECLIPSE Pour-On is rainfast. Rain occurring as early as 2 hours after treatment is unlikely to affect the efficacy of the product.

However rainfall on cattle less than 2 hours after treatment may result in reduced efficacy. ECLIPSE Pour-On should not be applied to wet animals. Ask your local veterinary practitioner or animal health advisor for recommended parasite management practices for your area to reduce development of resistance. It is advisable that a resistance test be conducted regularly when using any parasite treatment.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER THE APPROPRIATE LEGISLATION

WITHHOLDING PERIODS

MEAT: DO NOT USE less than 56 days before slaughter for human consumption. Calves MUST NOT be slaughtered for human consumption within 60 days of treatment of dams.

MILK: DO NOT USE in lactating cows or within 60 days of calving where milk may be used or processed for human consumption. If cows calve earlier than 60 days after treatment, milk may contain residues. This milk MUST NOT be used for human consumption, supplied for processing or fed to bobby calves for at least 60 days following treatment.

TRADE ADVICE

EXPORT SLAUGHTER INTERVAL (ESI)
DO NOT USE less than 70 days before slaughter for export. The ESI on this label was correct at the time of label approval.

Before using this product, confirm the current ESI from the manufacturer on 1800-808-691 or the APVMA website (www.apvma.gov.au/residues).

SAFETY DIRECTIONS

Harmful if swallowed or absorbed by skin contact. Repeated exposure may cause allergic disorders. Will damage the eyes and skin. Avoid contact with eyes and skin. If clothing becomes contaminated with product, remove clothing immediately. If product is in eyes, wash it out immediately with water.

When opening the container and using the product, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), a washable hat, elbow-length chemical resistant gloves and goggles.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If skin contact occurs, remove contaminated clothing and wash skin thoroughly.

ADDITIONAL SAFETY INFORMATION

When handling cattle for multiple procedures, it is recommended to apply ECLIPSE Pour-On last to minimise workers' exposure by contact with the treated animals. Additional information is listed in the Safety Data Sheet.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND THE ENVIRONMENT

Abamectin is extremely toxic to aquatic species. Do not contaminate dams, rivers, streams or other waterways with the chemical or used container.

DISPOSAL

This container can be recycled if it is clean, dry, free of visible residues and has the *drumMUSTER* logo visible. Triple or pressure rinse container for disposal. Dispose of rinseate or any undiluted chemical according to state legislative requirements. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any *drumMUSTER* collection or similar container management program site. The cap should not be replaced but may be taken separately.

STORAGE

Store below 25°C (Air Conditioning). Protect from light.

Merial Australia Pty Ltd.
Building D, 12-24 Talavera Road.
Macquarie Park, NSW, 2113

For product information and technical enquiries visit www.merial.com.au or call Merial Customer Solutions Australia: 1800 808 691. Made in New Zealand.

(B):

EXP:

APVMA Approval No. 60752/59249

