

H 30252 (AU)

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

ivomec[®]
Eprinex[™]
(eprinomectin)



Pour-On for Beef and Dairy Cattle

Active Constituent: EPRINOMECTIN 5 mg/mL

For the treatment and control of internal and external parasites of beef and dairy cattle and internal parasites of deer.

READ THE ENCLOSED LEAFLET BEFORE USING THIS PRODUCT.

Dose 1 mL per 10 kg liveweight. For external use only.

Merial Australia Pty Limited
A.C.N. 071 187 285
Level 6, 79 George Street
Parramatta NSW 2150

For product information and for
emergency advice call the 24-hour
toll-free Merial technical hotline
1800-808-691.

NRA Approval No. 49105/0198



IVOMEC and cattle head logo are registered trademarks and
EPRINEX is a trademark of Merial.

53381/300300C

2.5 Litre



ivomec®

Eprinex™

(eprinomectin)

Pour-On for Beef and Dairy Cattle

Active Constituent: EPRINOMECTIN 5 mg/mL

INDICATIONS:

For the treatment and control of gastrointestinal roundworms (including inhibited immature *Ostertagia ostertagi*) and lungworm in cattle.

IVOMECEPRINEX Pour-On for Beef and Dairy Cattle contains eprinomectin, a member of the macrocyclic lactone family of drenches. It is effective against eprinomectin sensitive strains of internal parasites. Resistance can develop to any drench.

For sucking and biting lice, sarcoptic and chorioptic mange mites and buffalo fly in beef and dairy cattle.

IVOMECEPRINEX Pour-On aids in control of cattle tick (*Boophilus microplus*). For the treatment and control of gastrointestinal roundworms and lungworm in deer.

PERSISTENT ACTIVITY IN CATTLE:

IVOMECEPRINEX Pour-On when given at the recommended dosage of 1 mL/10 kg liveweight controls infections with:

Parasite*	Prolonged Activity
<i>Ostertagia</i> spp.	Up to 28 days
<i>Cooperia</i> spp.	Up to 28 days
<i>Dictyocaulus viviparus</i>	28 days
<i>Nematodirus helvetianus</i>	Up to 28 days
<i>Oesophagostomum radiatum</i>	Up to 28 days
<i>Haemonchus placei</i>	21 days
<i>Trichostrongylus</i> spp.	21 days

*The following parasite species are included within each of the relevant genera: *Ostertagia ostertagi*, *O. leptospicularis*, *O. lyrata*, *Cooperia oncophora*, *C. punctata*, *C. surnabada*, *Trichostrongylus axei*, *T. colubriformis*.

DOSAGE AND ADMINISTRATION:

For pour-on application only. Apply along the topline in a narrow strip extending from the withers to the tailhead.

The dose rate is 1 mL for each 10 kg of liveweight.

WITHHOLDING PERIODS:

CATTLE MILK: NIL.

There is no milk withholding period for lactating dairy cattle.

CATTLE MEAT/DEER MEAT: NIL.

There is no meat withholding period for slaughter of cattle and deer. As the standards that are applied for eprinomectin in overseas markets are similar to those approved in Australia, the export slaughter interval (ESI) is the same as the withholding period i.e. EXPORT SLAUGHTER INTERVAL: NIL.

SAFETY:

Studies have demonstrated a wide safety margin. Five times the recommended use level had no adverse effect on cattle or deer. Three times the recommended use level had no adverse effect on breeding performance of cows and bulls.

NOTE TO USER:

Rainfall before or after treatment will not affect the efficacy of IVOMECEPRINEX Pour-On.

PRECAUTIONS:

Close container when not in use.

Do not apply to areas of the backline covered with mud or manure.

This product is for application to skin surface only, do not give orally or by injection.

ADMINISTRATION:

Connect the dosing gun to the pack as follows:

- Attach the open end of the draw-off tubing to an appropriate dosing applicator. Attach draw-off tubing to the cap with the stem that is included in the pack.
- Replace the shipping cap with the cap that has the draw-off tubing and tighten this cap.
- Gently prime the dosing applicator, checking for leaks.

Follow the dosing applicator manufacturer's directions for adjusting the dose and proper use and maintenance of the equipment.

STORAGE:

Store below 30°C (Room Temperature).

Store bottle or pack in carton to protect from light.

DISPOSAL:

Triple or (preferably) pressure rinse IVOMECEPRINEX Pour-On containers and dispose of rinsate in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Break, crush or puncture and bury empty containers at a local authority landfill. If not available, bury the containers below 500 mm in the disposal pit. Empty containers and product should not be burnt. Very highly toxic to aquatic species. Do not contaminate dams, rivers, streams or other waterways with the chemical or used container.

PROTECTION OF FISH, CRUSTACEANS AND THE ENVIRONMENT:

Studies indicate that when eprinomectin comes in contact with the soil, it readily and tightly binds to the soil and becomes inactive over time.

Containers and any residual contents should be disposed of safely in an approved landfill as free eprinomectin may adversely affect fish and certain water-borne organisms.

SAFETY DIRECTIONS:

Poisonous if swallowed.

Will irritate the eyes. Avoid contact with eyes. Repeated exposure may cause allergic disorders. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.

FIRST AID:

If poisoning occurs, contact a doctor or Poisons Information Centre, phone 131126. If swallowed, and if more than 15 minutes from a hospital, induce vomiting, preferably using Ipecac Syrup A.P.F.

For further information, refer to the Material Safety Data Sheet.

Batch Number:

Expiry Date:



53382/300300C

